

AD-A093 279

WISCONSIN UNIV-MADISON DEPT OF COMPUTER SCIENCES  
ON THE EIGENVALUES OF 2ND ORDER ELLIPTIC DIFFERENCE OPERATORS.(U)  
NOV 80 S V PARTER

F/6 12/1

N00014-76-C-0341

UNCLASSIFIED

CSTR-405

NL

[ 0 ]

8/1/80

END

DATE

FILED

1-8

DTIC

AD A093279


432 .

**LEVEL II**

**COMPUTER SCIENCES  
DEPARTMENT**

University of Wisconsin-  
Madison

(12)



Contract N00014-76-C-0341

ON THE EIGENVALUES OF 2ND  
ORDER ELLIPTIC DIFFERENCE OPERATORS

by

Seymour V. Parter

Computer Sciences Technical Report #405

November 1980

RECEIVED  
DEC 29 1980

FILE COPY

80 11 25 001

12

6

ON THE EIGENVALUES OF 2ND  
ORDER ELLIPTIC DIFFERENCE OPERATORS.

by

10) Seymour V. Parter

13) 12

9) Technical Repts.

11) Nov 28

DTIC  
ELECTED  
DEC 29 1980  
S C

14) ZSTR-4A

15) \*This work was supported by the Office of Naval Research under Contract No. N00014-76-C-0341.

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

7 37 124 216

## I. INTRODUCTION

Let  $L$  be a second order elliptic operator defined in a smooth domain  $\Omega \subset \mathbb{R}^n$ . That is

$$(1.1) \quad Lu = -\sum \frac{\partial}{\partial x_i} a_{ij}(x) \frac{\partial u}{\partial x_j} + \sum b_i(x) \frac{\partial u}{\partial x_i} + c(x)u$$

where, for every  $\xi \in \mathbb{R}^n$  we have

$$(1.2a) \quad \lambda_0 \sum \xi_j^2 \leq \sum a_{ij}(x) \xi_i \xi_j \leq \Lambda_0 \sum \xi_j^2, \quad \forall x \in \Omega,$$

for certain positive constants  $\lambda_0$ ,  $\Lambda_0$ , with

$$(1.2b) \quad 0 < \lambda_0 < \Lambda_0.$$

We may assume the coefficients  $a_{ij}(x)$ ,  $b_i(x)$ ,  $c(x) \in C^\infty(\bar{\Omega})$  and

$$(1.2c) \quad c(x) \geq 0.$$

Let  $N$  be a first order operator defined in  $\Omega$ , i.e.

$$(1.3) \quad Nu = \sum \beta_i(x) \frac{\partial u}{\partial x_i} + \alpha(x)u.$$

In this report we are concerned with the eigenvalue problem

$$(1.4a) \quad Lu = \lambda Nu \quad \text{in } \Omega$$

$$(1.4b) \quad u = 0 \quad \text{on } \partial\Omega$$

and the related discrete eigenvalue problem

$$(1.5a) \quad L_h u_h = \lambda_h N_h u_h \quad \text{in } \Omega(h)$$

$$(1.5b) \quad u_h = 0 \quad \text{on } \partial\Omega(h)$$

Accession For	<input checked="" type="checkbox"/>
NTIS GRA&I	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By	
Distribution	
Availability	
Distribution	
Dist	

A

where  $L_h$  and  $N_h$  are finite-difference operators defined on a grid region  $\Omega(h)$  with boundary  $\partial\Omega(h)$ .

In the case where  $L$  and  $N$  are self-adjoint, i.e.

$$(1.6) \quad b_i(x) = 0, \quad \beta_i(x) = 0, \quad i = 1, 2, \dots, n$$

and  $\alpha(x)$  does not change sign in  $\Omega$  while  $L_h$  and  $N_h$  correspond to positive definite matrices one can attack this problem via the appropriate variational principles (see Weinberger [12]).

In the general case if  $L_h$  and  $N_h$  arise from a Galerkin approach and one has general  $L_2(\Omega)$  convergence results one may apply the results of Vainikko [10] and Osborn [7].

At first glance it would appear that one could somehow easily modify the finite-difference set-up so as to be able to apply the finite-element theory. This author was unable to do this.

The difficulty centers about the following points. Let

$$T = L^{-1}N, \quad T_h = L_h^{-1}N_h.$$

In order to apply the theory of [7] we require that

$$T_h f \rightarrow T f \quad \forall f \in X$$

where  $X$  is an appropriate Banach space, say  $L_2(\Omega)$ . But, the finite-difference theory is not designed to operate on all of  $L_2(\Omega)$ , only on sufficiently smooth functions.

Moreover, if one attempts to extend the theory to all of  $L_2(\Omega)$  via  $L_2$  projection or "smoothing", one finds that the eigenvalue problem be-

comes perturbed in a manner which is non-trivial when viewed from all of  $L_2(\Omega)$ . On the other hand, a glance at the development of [7] shows that all of the "real action" takes place in certain smooth, finite dimensional subspaces; the eigenspaces. This remark is the key to the development given in this report.

→ This work is motivated by the desire to extend the theory of iterative methods for the solution of the algebraic problems associated with  $L_p^{subh}$  de, a finite difference operator. developed in [8] to the non-self-adjoint case. See [9]. → For this reason the operators  $N_k^{subh}$  will approximate  $N$  in a relatively "weak" manner.

In order to simplify the presentation we first consider a particular "model" problem. Following that development we discuss the essential features of that discussion. ✓

In section 2 we describe the basic model problem and some well-known facts about this model problem.

(In section 3 we develop the convergence theorems.

In section 4 we discuss the ideas in our convergence proofs and their possible extension to more general problems. ↗

## 2. THE MODEL PROBLEM

Let

$$(2.1) \quad \Omega \equiv \{(x,y); 0 < x, y < 1\}$$

and let

$$(2.2) \quad Lu = -(u_{xx} + u_{yy}) + a(x,y)u_x + b(x,y)u_y$$

where  $a, b \in C^\infty(\overline{\Omega})$  are nice functions. Let

$$(2.3) \quad Nu = \alpha(x,y)u_x + \beta(x,y)u_y + q(x,y)u$$

with  $\alpha, \beta, q \in C^\infty(\overline{\Omega})$ . For every  $f \in L_2(\Omega)$  we let

$$Tf = u$$

be the solution of

$$(2.4a) \quad Lu = Nf, \text{ in } \Omega,$$

$$(2.4b) \quad u = 0, \text{ on } \partial\Omega.$$

Since  $N$  maps  $L_2(\Omega)$  into  $H_{-1}(\Omega)$  and the resulting  $u$  is in  $H_1(\Omega)$  we see that  $T$  is a compact map whose spectrum  $\sigma(T)$  consists only of eigenvalues and 0. Thus a point  $z \in \mathbb{C}$ ,  $z \neq 0$ , is in the spectrum of  $T$  only if there is a  $u \in L_2(\Omega)$  which satisfies

$$(2.5) \quad Tu = zu, \quad u \neq 0.$$

We assume that 0 is not an eigenvalue of  $T$ .

Let  $P$  be a fixed positive integer and set

$$h = \frac{1}{P+1} .$$

Let

$$(2.6a) \quad \Omega(h) \equiv \{(x_i, y_j) = (ih, jh); 1 \leq i, j \leq P\}$$

be the set of interior mesh points while

$$(2.6b) \quad \partial\Omega(h) \equiv \{(x_i, y_j) = (ih, jh); i \text{ or } j = 0 \text{ or } P+1\}$$

is the set of boundary mesh points. A "grid vector"  $\{U_{ij}; 0 \leq i, j \leq P+1\}$  is a function defined on the total set of mesh points

$$\bar{\Omega}(h) = \Omega(h) \cup \partial\Omega(h) .$$

As usual, we define the discrete Laplace Operator by: for  $1 \leq i, j \leq P$

$$(2.7) \quad [\Delta_h U]_{ij} = \frac{U_{i+1,j} - 2U_{i,j} + U_{i-1,j}}{h^2} + \frac{U_{i,j+1} - 2U_{i,j} + U_{i,j-1}}{h^2} .$$

Let

$$(2.8a) \quad a_{ij} = a(x_i, y_i) , \quad b_{ij} = b(x_i, y_i) .$$

With this notation we now define the finite-difference operator  $L_h$  corresponding to  $L$ . For  $1 \leq i, j \leq P$

$$(2.8) \quad [L_h U]_{ij} = -[\Delta_h U]_{ij} + a_{ij} \frac{U_{i+1,j} - U_{i-1,j}}{2h} + b_{ij} \frac{U_{i,j+1} - U_{i,j-1}}{2h} .$$

Remark: While  $U$  is defined on the entire mesh region, the values  $\Delta_h U$  and  $L_h U$  are defined only on the interior mesh points,  $\Omega(h)$ .



We also define the usual difference operators

$$(2.9a) \quad [\nabla_x U]_{i,j} = \frac{U_{i+1,j} - U_{i,j}}{h}, \quad 1 \leq j \leq P, \quad 0 \leq i \leq P$$

$$(2.9b) \quad [\nabla_y U]_{i,j} = \frac{U_{i,j+1} - U_{i,j}}{h}, \quad 0 \leq j \leq P, \quad 1 \leq i \leq P.$$

Let  $U, V$  be grid vectors. We define an inner product

$$(2.10a) \quad (U, V) = h^2 \sum_{i=1}^P \sum_{j=1}^P \bar{U}_{ij} V_{ij}.$$

Observe that this inner product involves only the interior mesh points,  
 $\Omega(h)$ . In the usual way we define the semi-norm

$$(2.10b) \quad \|U\|_h^2 = (U, U)$$

We note that this semi-norm is a norm for those grid vectors  $U$  which vanish on the boundary, i.e.,

$$(2.11) \quad U = 0 \quad \text{on} \quad \partial\Omega(h).$$

We recall the following basic facts.

Lemma 2.1: Let  $U$  be a grid vector which satisfies (2.11). Then

$$(2.12) \quad (-\Delta_h U, U) = (\nabla_x U, \nabla_x U) + (\nabla_y U, \nabla_y U).$$

Proof: Apply summation by parts. See [1], [11].

Definition: Let  $f(x, y) \in C(\bar{\Omega})$ . Then  $Q_h f$  is the grid vector which interpolates at the mesh points. That is,  $[Q_h f]$  is the restriction of  $f$  to  $\Omega(h)$ ;

$$(2.13) \quad [Q_h f]_{ij} = f_{ij} = f(x_i, y_j).$$

Definition: Let  $f \in C^2(\bar{\Omega})$  and let  $\omega_2(f;h)$  be the modulus of continuity of the second derivatives of  $f$ , i.e. Let  $\phi(x,y)$  be any second derivative of  $f$ ,  $\phi = f_{xx}$  or  $\phi = f_{yy}$  or  $\phi = f_{xy}$ . For all  $(x,y) \in \bar{\Omega}$  and  $(x',y') \in \bar{\Omega}$  with

$$|x-x'| \leq h, \quad |y-y'| \leq h,$$

we have

$$(2.14) \quad \sup \{ |\phi(x,y) - \phi(x',y')|; \phi = f_{xx}, f_{xy}, f_{yy} \} = \omega_2(f;h).$$

Of course,  $\omega_2(f,h) \rightarrow 0$  as  $h \rightarrow 0$ .

Definition: Let  $L^*$  be the formal adjoint of the operator  $L$ . That is

$$(2.15a) \quad L^*U = -(U_{xx} + U_{yy}) - (au)_x - (bu)_y.$$

Let  $L_h^*$  be the finite-difference operator defined by

$$(2.15b) \quad [L_h^*U]_{ij} = -[\Delta_h U]_{ij} - \left[ \frac{a_{i+1,j}U_{i+1,j} - a_{i-1,j}U_{i-1,j}}{2h} \right] - \left[ \frac{b_{i,j+1}U_{i,j+1} - b_{i,j-1}U_{i,j-1}}{2h} \right].$$

Lemma 2.2: Let  $U, V$  be grid vectors which vanish on  $\partial\Omega(h)$ , i.e. both  $U$  and  $V$  satisfy (2.11). Then

$$(2.16) \quad (U, L_h V) = (L_h^* U, V).$$

If  $U \in C^2(\bar{\Omega})$  then there is a constant  $K > 0$  such that

$$(2.17a) \quad L_h Q_h U = Q_h L U + \tau$$

$$(2.17b) \quad L_h^* Q_h U = Q_h L^* U + \tau^*$$

where

$$(2.17c) \quad \|\tau\|_\infty + \|\tau^*\|_\infty \leq K \omega_2(U, h) .$$

Proof: The identity (2.16) follows from an elementary summation by parts.

The estimate (2.17c) follows from Taylor's theorem. Of course, if  $U \in C^4(\bar{\Omega})$  then the truncation error  $|E| + |E^*|$  is of order  $h^2$ .

Lemma 2.3: Let  $F$  be a grid vector and let

$$\|F\|_\infty = \max \{|F_{ij}|; 1 \leq i, j \leq P\} .$$

There are constants  $K > 0$ ,  $h_0 > 0$  such that, if

$$0 < h \leq h_0$$

and  $U$  is the solution of

$$(2.18a) \quad L_h U = F \quad \text{in } \Omega(h) ,$$

$$(2.18b) \quad U = 0 \quad \text{on } \partial\Omega(h) ,$$

then

$$(2.19) \quad \|U\|_\infty \leq K \|F\|_\infty .$$

Proof: This well-known result follows from the discrete maximum principle and lemma 2.2. See [11].

Lemma 2.4: Let  $U$  be a grid vector which vanishes on  $\partial\Omega(h)$ . Then

$$(2.20) \quad -(\Delta_h U, U) \leq |(U, L_h U)| + \|a\|_\infty \|U\|_h \cdot \|\nabla_x U\|_h \\ + \|b\|_\infty \|U\|_h \cdot \|\nabla_y U\|_h .$$


Proof: Direct Computation.

Before discussing the operators  $N_h$  we discuss the "prolongation" of grid vectors  $U$  into functions  $U(x,y;h)$  defined on  $\Omega$ .

Let the mesh region  $\bar{\Omega}(h)$  be "triangulated". To be specific, imagine that the diagonal lines

$$y = x - jh, \quad j = -(P+1), -P, \dots, P, (P+1)$$

have been drawn over the grid. Each square with vertices  $(x_i, y_j)$ ,  $(x_{i+1}, y_j)$ ,  $(x_{i+1}, y_{j+1})$ ,  $(x_i, y_{j+1})$  is considered as the union of two triangles.



With each grid vector  $U = \{U_{ij}\}$  we associate a piecewise linear function  $U(x,y;h)$  which is linear on each triangle and

$$U(x_i, y_j, h) = U_{ij} .$$

Let  $P(h)$  be the space of all such function and  $P_0(h)$  be the subspace of  $P(h)$  which consists of those functions which vanish on  $\partial\Omega$ .

Throughout the remainder of this work we will make a complete identification between grid vectors  $U = \{U_{ij}\}$  and these associated functions  $U(x,y;h)$ . This will avoid cumbersome notational difficulties.

Let  $\langle \cdot, \cdot \rangle$  be the  $L_2(\Omega)$  inner product, i.e. if  $f, g \in L_2(\Omega)$ , then

$$(2.21) \quad \langle f, g \rangle = \int_{\Omega} \int \bar{f}(x,y)g(x,y)dxdy.$$

Lemma 2.5: Let  $U, V \in P_0(h)$ . Then

$$(2.22) \quad \begin{aligned} \langle U, V \rangle &= \frac{10}{24} \langle U, V \rangle + \\ &\quad \frac{3h^2}{24} \sum_{i,j} \bar{U}_{ij} (V_{i+1,j} + V_{i-1,j} + V_{i,j+1} + V_{i,j-1}) \\ &\quad + \frac{h^2}{24} \sum_{i,j} \bar{U}_{ij} (V_{i+1,j-1} + V_{i-1,j+1}). \end{aligned}$$

Furthermore

$$(2.23a) \quad \langle U, V \rangle = (U, V) + E_1$$

where

$$(2.23b) \quad |E_1| \leq h(U, U)^{\frac{1}{2}} (-\Delta_h V, V)^{\frac{1}{2}}.$$

Proof: The formula (2.22) is obtained by a direct integration. The estimates (2.23a) and (2.23b) follow from (2.22) and lemma 2.1.

Lemma 2.6: Let  $U \in P(h)$ . Let  $S_h$  be the triangle with vertices  $I, J, K$ . Then

$$(2.24) \quad \int_{S_h} U(x, y; h) dx dy = \frac{h^2}{6} (U_I + U_J + U_K) .$$

In this notation  $I, J$  and  $K$  are symbols for pairs of indices  $(i, j)$ .

Proof: Direct integration.

We now turn to a discussion of the operator  $N_h$ . We do not describe these operators as specifically as we described  $L_h$  and  $L_h^*$ . Instead, we make the following weak consistency assumption.

Assumption N: Let  $N_h^0$  be defined by: for  $1 \leq i, j \leq P$

$$\begin{aligned} [N_h^0 U]_{ij} = & \alpha(x_i, y_j) \left[ \frac{U_{i+1,j} - U_{i-1,j}}{2h} \right] + \beta(x_i, y_j) \left[ \frac{U_{i,j+1} - U_{i,j-1}}{2h} \right] \\ & + q(x_i, y_j) U_{ij} . \end{aligned}$$

We assume that, for all grid vectors  $U, V$  which vanish on  $\partial\Omega(h)$ ,

$$(2.25a) \quad (U, N_h V) = (U, N_h^0 V) + E_2$$

where

$$\begin{aligned} (2.25b) \quad |E_2| \leq & Dh [(-\Delta_h U, U) \cdot (-\Delta_h V, V)]^{\frac{1}{2}} \\ & + Dh^2 [(-\Delta_h U, U) + (-\Delta_h V, V)] \end{aligned}$$

Lemma 2.7: Let  $U \in P_0(h)$ . Then there is a constant  $C_1$ , depending on  $\|\alpha\|_\infty, \|\beta\|_\infty, \|q\|_\infty$ , but not on  $h$  so that

$$(2.26a) \quad (\overset{0}{N}_h U, \overset{0}{N}_h U) \leq C_1 (-\Delta_h U, U) .$$

$$(2.26b) \quad |(V, \overset{0}{N}_h U)| \leq C_1 \|V\|_h (-\Delta_h U, U)^{\frac{1}{2}} .$$

Proof: Direct Computation.

Before proceeding to the convergence theorem of the next section, we recall some basic facts.

If  $U \in P_0(h)$  then

$$(2.27) \quad (-\Delta_h U, U) = \|U\|_{\overset{0}{H}_1(\Omega)}^2$$

Moreover, we have the important

Lemma 2.8 (Rellich): Let  $h_n \rightarrow 0$  and let  $U_n \in P_0(h_n)$  satisfy

$$(-\Delta_{h_n} U_n, U_n) = \|U_n\|_{\overset{0}{H}_1(\Omega)}^2 \leq K .$$

Then there is a subsequence  $n'$  and a function  $u \in \overset{0}{H}_1(\Omega)$  such that

$$U_{n'} \rightharpoonup u \text{ in } \overset{0}{H}_1(\Omega)$$

$$U_{n'} \rightarrow u \text{ in } L_2(\Omega) .$$

Proof: See [2].

### 3. THE CONVERGENCE THEOREMS

Lemma 3.1: Let  $\Sigma \subset \mathbb{C}$  be a compact subset of the resolvent set of  $T$ .

Assume that

$$0 \notin \Sigma.$$

For each  $z \in \Sigma$  consider the problem

$$(3.1a) \quad L_h U - \frac{1}{z} N_h U = F, \quad \text{in } \Omega(h),$$

$$(3.1b) \quad U = 0, \quad \text{on } \partial\Omega(h).$$

There are constants  $h_1 > 0$ ,  $\bar{K} > 0$  with  $\bar{K}$  independent of  $h$ , such that: For  $0 < h \leq h_1$ , problem 3.1 has a unique solution  $U$ , and

$$(3.2) \quad \|U\|_h \leq \bar{K} \|F\|_h.$$

Moreover, for every  $u \in C^2(\bar{\Omega})$  which vanishes on  $\partial\Omega$ , let  $V$  be the grid vector which satisfies

$$(3.3a) \quad L_h V - \frac{1}{z} N_h V = Q_h [Lu - \frac{1}{z} Nu], \quad \text{in } \Omega(h)$$

$$(3.3b) \quad V = 0, \quad \text{on } \partial\Omega(h).$$

Then

$$\langle V-u, V-u \rangle \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

That is,  $V(x,y;h)$  converges to  $u$  in  $L_2(\Omega)$ .



Proof: It is only necessary to verify (3.2). Suppose (3.2) is false. Then there are sequences  $z_n \in \Sigma$ ,  $h_n \rightarrow 0$  and grid vectors  $F_n$  such that

$$(3.4a) \quad \|F_n\|_{h_n} \rightarrow 0 \text{ as } h_n \rightarrow 0,$$

while the corresponding solutions [(3.1a), (3.1b)],  $V_n$ , satisfy

$$(3.4b) \quad \|V_n\|_{h_n} = 1.$$

Then

$$(3.5a) \quad L_{h_n} V_n - \frac{1}{z_n} N_{h_n} V_n = f_n, \quad \text{in } \Omega(h_n),$$

$$(3.5b) \quad V_n = 0, \quad \text{on } \partial\Omega(h_n).$$

$$(3.5c) \quad \|V_n\|_{h_n} = 1.$$

Applying lemma 2.4 we have

$$(3.6) \quad \begin{aligned} -(\Delta_{h_n} V_n, V_n) &\leq |(V_n, f_n)| + \|a\|_\infty \|\nabla_x V_n\|_{h_n} + \|b\|_\infty \|\nabla_y V_n\|_{h_n} \\ &\quad + \frac{1}{|z_n|} |(V_n, N_{h_n} V_n)|. \end{aligned}$$

Using Assumption N and Lemma 2.7 we have

$$|(V_n, N_{h_n} V_n)| \leq C_1 (-\Delta_{h_n} V_n, V_n)^{\frac{1}{2}} + 2Dh_n (-\Delta_{h_n} V_n, V_n)$$

Therefore

$$(3.7) \quad -(\Delta_{h_n} V_n, V_n) \leq \|f_n\|_{h_n} + K_2 + \left(\frac{1}{4} + Dh_n\right) (-\Delta_{h_n} V_n, V_n)$$

where

$$K_2 = 10(\|a\|_\infty^2 + \|b\|_\infty^2 + \bar{c}_1^2)$$

$$\bar{c}_1 = \max \left\{ \frac{C_1}{|z|}, z \in \Sigma \right\}.$$

$$\bar{D} = \max \left\{ \frac{2D}{|z|}, z \in \Sigma \right\}.$$

Thus, when

$$\bar{D}h_n \leq \frac{1}{4}$$

we obtain

$$(3.8) \quad (-\Delta_{h_n} V_n, V_n) \leq 2(1 + K_2).$$

Let  $z_{n_i}$  be a subsequence of the  $z_n$  such that

$$z_{n_i} \rightarrow \zeta \in \Sigma.$$

Applying Lemma 2.8 we see that there is a sub-subsequence  $n''$  and a function  $v \in H_1^0$  such that

$$(i) \quad V_{n''} \rightarrow v \text{ in } H_1^0(\Omega)$$

and

$$(ii) \quad V_{n''} \rightarrow v \text{ in } L_2(\Omega).$$

For simplicity of notation we denote  $\{h_{n''}\}$  by  $\{h_n\}$ .

Let  $\phi \in C_0^\infty(\Omega)$  and let

$$(3.9) \quad \phi_n = Q_{h_n} \phi.$$

From (3.5a), (3.5b) we obtain

$$(\phi_n, L_{h_n} V_n) - \frac{1}{z_n} (\phi_n, N_{h_n} V_n) = (\phi_n, f_n) .$$

Thus, using Assumption N we see that

$$(L_{h_n}^* \phi_n, V_n) - \frac{1}{z_n} (\phi_n, N_{h_n}^0 V_n) = (\phi_n, g_n)$$

where

$$\|g_n\|_{h_n} \rightarrow 0 \text{ as } n \rightarrow \infty .$$

On letting  $n \rightarrow \infty$  lemma 2.2 and lemma 2.5 yield

$$(3.10) \quad \langle L^* \phi, v \rangle - \frac{1}{\zeta} \langle \phi, N v \rangle = 0 .$$

Thus,  $v$  is a weak solution of

$$(3.11a) \quad L v - \frac{1}{\zeta} N v = 0 , \text{ in } \Omega ,$$

$$(3.11b) \quad v = 0 , \text{ on } \partial \Omega .$$

But, then,  $v$  is a genuine solution of (3.11a), (3.11b) and  $\zeta \in \Sigma$  is an eigenvalue of  $T$  (see [2]). With this contradiction we have proven (3.2).

The remainder of the lemma follows from similar arguments.

Let  $\mu \neq 0$  be an eigenvalue of  $T$ . Let

$$S \equiv \ker (\mu - T)^\alpha ,$$

i.e., the kernel or nullspace of the linear map  $(\mu - T)^\alpha$  where  $\alpha$  is the ascent or rank of  $\mu$  (see [5]).  $S$  is the range of the linear

projection  $E$  given by

$$(3.12) \quad E = \frac{1}{2\pi i} \int_{\Gamma} (z-T)^{-1} dz$$

with  $\Gamma$  any circle in the resolvent set of  $T$  enclosing  $\mu$  in its interior, but no other point of the spectrum. The linear projector  $E$  can also be written in the form

$$(3.13a) \quad Ey = \sum_{i=1}^a \langle y, \phi_i^* \rangle \phi_i$$

where  $\{\phi_i, i=1,2,\dots,a\}$  is any basis for  $S$  and  $\{\phi_i^*, i=1,2,\dots,a\}$  is the corresponding dual basis in

$$(3.13b) \quad S^* = \ker (\mu - T^*)^\alpha .$$

That is  $\phi_i^* \in S^*$  and  $\langle \phi_i, \phi_j^* \rangle = \delta_{ij}$ . Here  $T^*$  is determined by

$$(3.13c) \quad \langle Tu, v \rangle = \langle u, T^*v \rangle .$$

That is,

$$(3.13d) \quad T^* = N^*(L^*)^{-1} .$$

Let  $\Sigma = \Gamma$ . Applying Lemma 3.1 we see that

$$(3.14) \quad (z - T_h)^{-1} = \frac{1}{z} (L_h - \frac{1}{z} N_h)^{-1} L_h$$

is well-defined on  $\Gamma$  for  $0 < h \leq h_1$ . Let

$$(3.15) \quad E_h = \frac{1}{2\pi i} \int_{\Gamma} (z - T_h)^{-1} dz .$$

Lemma 3.2: Let  $\mu \neq 0$  be an eigenvalue of  $T$ . There is an  $h_2 > 0$  such that if  $0 < h \leq h_2$  there is an eigenvalue  $\mu_h$  of  $T_h$  which lies in the interior of  $\Gamma$ .

Proof: If the lemma is false, then

$$E_h \equiv 0.$$

Let  $\phi \in S$  be an eigenfunction of  $T$  with eigenvalue  $\mu$ . It is not difficult to prove that  $\phi \in C^2(\overline{\Omega})$ . However, in order to see how the general argument goes, we replace  $\phi$  by a smooth  $H_2^0 \cap H_1^0$  approximant, which we again call  $\phi$ . Let

$$(3.16a) \quad \phi_h = Q_h \phi, \quad \psi_h = E_h \phi_h$$

and let

$$(3.16b) \quad \psi_h = Q_h E \phi,$$

we observe that

$$\psi_h(x, y; h) \rightarrow E \phi \neq 0.$$

We calculate

$$\psi_h - \psi_h = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z} [Q_h (L - \frac{1}{z} N)^{-1} L \phi - (L_h - \frac{1}{z} N_h)^{-1} L_h \phi_h].$$

Since  $\phi$  is smooth we may apply lemma 2.2. Thus

$$L_h \phi_h = Q_h L \phi + \tau$$

where  $\|\tau\|_\infty \rightarrow 0$ . In fact, for each fixed  $z \in \Gamma$  let  $v(x,y;z)$  be the solution of

$$(3.17a) \quad (L - \frac{1}{z}N)v = L\phi, \quad v = 0 \text{ on } \partial\Omega,$$

while  $V$  is the solution of

$$(3.17b) \quad (L_h - \frac{1}{z}N_h)V = Q_h L_h \phi, \quad V = 0 \text{ on } \partial\Omega(h).$$

From Lemma 3.1 we see that

$$\|V\|_h + (-\Delta_h V, V)^{\frac{1}{2}} \leq K \|L\phi\|_\infty.$$

Thus, using Lemma 2.5 and Hölder's inequality we see that there is a constant  $K_1$  such that

$$(3.18) \quad \|V\|_{L_2(\Omega)} \leq K_1, \quad \|V\|_{L_1(\Omega)} \leq K_1.$$

Thus

$$(3.19) \quad u(x,y;h) - V(x,y;h) = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z} (v(x,y;z) - V(x,y;z)) dz.$$

Applying Lemma 3.1 we see that

$$\|u - V\|_{L^2(\Omega)} \leq \frac{1}{2\pi} \int_0^{2\pi} \theta d\theta \|v(\cdot, \cdot; z) - V(\cdot, \cdot; z)\|_{L^2(\Omega)}.$$

However, according to Lemma 3.1 the integrand goes to 0 for every  $z \in \Gamma$  and the estimates (3.18) allow us to apply the dominated convergence theorem.

Thus

$$V(x,y;h) \rightarrow E\phi \neq 0.$$

Corollary: Let  $S_h$  denote the range of  $E_h$ . Then, for  $0 < h \leq h_2$  we have

$$\dim S_h \geq \dim S = a.$$

Proof: We need only apply the above argument to each  $\phi_i$  in the basis of  $S$ , smoothed if necessary. The resulting grid vectors  $E_h Q_h \phi_i$  will be linearly independent if  $h$  is small enough.

Lemma 3.3:  $\dim S = \dim S_h$ .

Proof: Let  $\mu_h \neq 0$  and  $V_h$  be an eigen pair of  $T_h$ . That is,

$$L_h V_h = \frac{1}{\mu_h} N_h V_h.$$

Suppose  $\mu_h \rightarrow \bar{\mu}$ . Then, it follows from Lemma 3.1 that  $\bar{\mu}$  is an eigenvalue of  $T$ .

In fact, let  $V_h$  satisfy

$$(\mu_h - T_h)^{r-1} V_h = W_h \neq 0$$

$$(\mu_h - T_h)^r V_h = 0.$$

A modification of the argument of Lemma 3.1 shows that, after selection of a subsequence if necessary, there are functions  $v, w \in L_2(\Omega)$  such that

$$V_h \rightarrow v, \quad W_h \rightarrow w \neq 0$$

and

$$(\bar{\mu} - T)^{r-1} v = w \neq 0$$

$$(\bar{\mu} - T)^r v = (\bar{\mu} - T)w = 0.$$

Thus, a basis for  $S_h$  converges to a set of linearly independent functions in  $S$ . Therefore

$$\limsup \dim S_h \leq \dim S = a.$$

Thus, the lemma follows from the corollary to Lemma 3.2.

We collect our results in the following

**Theorem 3.1:** Let  $\mu \neq 0$  be an eigenvalue of  $T$  with algebraic multiplicity  $a$ . Let  $\Gamma$  be a circle about  $\mu$  which does not include any other point of the spectrum of  $T$  either in its interior or on its boundary. Then there is an  $\bar{h}_1 > 0$  such that if  $0 < h \leq \bar{h}_1$

- (i) there are no eigenvalues  $\mu_h$  of  $T_h$  on  $\Gamma$
- (ii) there are eigenvalues  $\mu_{1,h}, \mu_{2,h}, \dots, \mu_{r,h}$  of  $T_h$  which lie inside  $\Gamma$ .
- (iii) Let  $m_1, m_2, \dots, m_r$  be the algebraic multiplicities of  $\mu_{1,h}, \mu_{2,h}, \dots, \mu_{r,h}$  respectively.

Then

$$\sum_{j=1}^r m_j = a.$$

Furthermore, the mapping

$$J_h = E_h Q_h|_S$$

is one to one and onto from  $S$  to  $S_h$ . Similarly, let

$$\tilde{J}_h = E|_{S_h}.$$



Then  $\tilde{J}_h$  is also one to one and onto from  $\tilde{S}_h$  to  $S$ .

Once one has this result one can follow Kreiss [6], Osborn [7] or de Boor and Swartz [3] to obtain Theorem 3.2: Let

$$\bar{\mu} = \frac{1}{a} \sum_{(j)} m_j \mu_{j,h}.$$

Then there is a constant  $K$  such that

$$|\bar{\mu} - \mu| \leq K \sup (T - T_h)|_S.$$

Specifically, we have

Let  $\phi_i$ ,  $i=1,2,\dots,a$  be a basis for  $S$  and  $\phi_i^*$ ,  $i=1,2,\dots,a$  be a basis for  $S^*$ . Then there is a constant  $K$  such that

$$|\bar{\mu} - \mu| \leq K \sum_{\sigma=1}^a \langle (T - T_h)\phi_\sigma, \phi_\sigma^* \rangle$$

Proof: See the development in [3].

#### 4. THE GENERAL IDEA

While the discussion in section 3 is limited to the model problem described in section 2, one can see the essential points and describe the basic ideas which will enable one to discuss more general problems.

The basic tool is the relationship between the linear projectors  $E$  given by (3.12) and  $E_h$  given by (3.15).

A look at the proof of Lemma 3.2 shows that one need only apply  $E_h$  to "smooth" elements  $\phi_h$ . However, even when we apply  $E_h$  to a smooth  $\phi_h$ , we must have results such as those given in Lemma 3.1.

Thus it is worthwhile to study the argument in Lemma 3.1. Almost any discussion of the operator  $L_h$  yields estimates such as those given in Lemma 2.2 (consistency) and Lemma 2.3 (convergence in the smooth case). However, the use of  $E$  and  $E_h$  requires estimates on the solutions  $V_h$  of the equations

$$zL_h V - N_h V = f_h, \quad z \in \Gamma.$$

We have essentially used estimates on  $L_h^{-1}$  to obtain these estimates. The basic facts are

FACT 1: If  $\{V_h; 0 < h \leq h_0\}$  is the set of all "functions" in  $P(h)$  such that

$$\|L_h V_h\|_h \leq C, \quad 0 < h \leq h_0,$$

then the  $\{V_h\}$  are sequentially compact.

FACT 2: If  $f_h, V_h \in P(h)$  and

$$(4.1a) \quad zL_h V_h - N_h^0 V_h = f_h$$

$$(4.1b) \quad f_h \rightarrow f \in C^2$$

$$(4.1c) \quad V_h \rightarrow V$$

then, in an appropriate sense

$$(4.2) \quad zLV - NV = f.$$

In the proof of Lemma 3.1 we first use these facts to establish (3.2). Then we use these facts and (3.2) to prove the convergence of the solutions of (3.3a), (3.3b).

Thus, in the more general case, i.e. general  $L, N$  and  $\Omega$ , one would expect to prove analogous spectral convergence theorems if one can establish these two facts. And, our discussion shows that this is the case. The proof of these facts are, in general, not too difficult. We have chosen to use "energy estimates" to establish FACT 1 and Weyl's lemma (weak solutions are genuine solution) to establish FACT 2.

In any particular instance one has a variety of possible arguments which lead to such conclusions. For example, in the case discussed in this report, i.e. the model problem, one could use the estimates of A. Brandt [4] to establish FACT 1. In fact, in this case, one can use the Banach space  $C(\Omega)$  rather than  $L_2(\Omega)$  and rely completely on the estimates of [4]. We choose to use "energy estimates" because we feel that they are (i) weaker and (ii) (generally) easier to obtain.

## REFERENCES

- [1] Aziz, A. K. and B. E. Hubbard: Bounds for the solution of the Sturm-Liouville problem with applications to finite difference methods, J. SIAM 12, 163-178 (1964).
- [2] Bers, L. F. John and M. Schechter: Partial Differential Equations, interscience N.Y., 1964.
- [3] de boor, C. and B. Swartz: Collocation Approximation to Eigenvalues of an Ordinary Differential Equation: The Principle of the Thing. Univ. of Wis. Math. Research Center Report 1937, (1979).
- [4] Brandt, A.: Estimates for difference quotients of solutions of Poisson type difference equations, Math. Comp. 20, 473-400 (1966).
- [5] Kato, T.: Perturbation Theory for Linear Operators, Grundlehren der Math. Wissenschaften, Band 132, Springer-Verlag, New York, 1966.
- [6] Kreiss, H.O.: Difference Approximations for Boundary and Eigenvalue Problems for Ordinary Differential Equations. Math. Comp. 26, 605-624 (1972).
- [7] Osborn, John, E.: Spectral Approximation for Compact Operators, Math. Comp. 29, 712-725 (1975).
- [8] Parter, S. V. and M. Steuerwalt: Another Look at Iterative Methods for Elliptic Difference Equations. Univ. of Wis. Computer Sciences Dept. Report #358 (1979).
- [9] Parter, S. V. and M. Steuerwalt: Report in Preparation
- [10] Vainikko, G. M.: Rapidity of Convergence of Approximation Methods in the Eigenvalue Problem. Ž. Vyčisl. Mat. i Mat. Fiz 7, 977-987 (1967) = U.S.S.R. Comput. Math. and Math. Phys. 7 18-32 (1967).

- [11] Varga, R. S.: Matrix Iterative Analysis, Prentice-Hall, Englewood Cliffs, N.J. (1962).
- [12] Weinberger, Hans F.: Variational Methods for Eigenvalue Approximation, Regional Conference Series in Applied Mathematics, No. 15, published by SIAM, Philadelphia, Penn. 1974.